

(19) World Intellectual Property Organization  
International Bureau




(43) International Publication Date  
17 April 2003 (17.04.2003)

PCT

(10) International Publication Number  
**WO 03/031587 A2**

- (51) International Patent Classification<sup>7</sup>: **C12N**
- (21) International Application Number: **PCT/US02/32503**
- (22) International Filing Date: 9 October 2002 (09.10.2002)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:  
60/328,162 9 October 2001 (09.10.2001) US  
60/328,689 10 October 2001 (10.10.2001) US
- (71) Applicant (for all designated States except US): **THE REGENTS OF THE UNIVERSITY OF CALIFORNIA** [US/US]; 1111 Franklin Street, 12th Floor, Oakland, CA 94607-5200 (US).
- (71) Applicants and  
(72) Inventors: **CARSON, Dennis, A.** [US/US]; 14824 Vista del Oceano, Del Mar, CA 92014 (US). **COT-TAM, Howard, B.** [US/US]; 3144 Willow Creek Place, Escondido, CA 92027 (US). **LEONI, Lorenzo, M.** [CH/US]; 3908 Via Tranquilo, San Diego, CA 92122 (US). **BARCHECHATH, Sylvie** [US/US]; 9321 Discovery Way, La Jolla, CA 92037 (US).
- (74) Agents: **STEFFEY, Charles, E.** et al.; Schwegman, Lundberg, Woessner & Kluth, P.O. Box 2938, Minneapolis, MN 55402 (US).
- (81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.
- (84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).
- Published:**  
— without international search report and to be republished upon receipt of that report
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: USE OF STAT-6 INHIBITORS AS THERAPEUTIC AGENTS



(57) Abstract: The present invention provides A therapeutic method to enhance the efficacy of interferon treatment comprising administering to a mammal subject to interferon treatment a compound which is an antagonist of the IL-4 or IL-13 signal transduction pathway in an amount effective to enhance said efficacy. The method includes treatment of diseases such as cancer, proliferative fibrotic diseases, viral diseases, or autoimmune diseases. The invention also includes the use of chemotherapeutic agents, radiation or other treatments in conjunction with the method of the invention.

## USE OF STAT-6 INHIBITORS AS THERAPEUTIC AGENTS

5

### Cross-Reference to Related Applications

This application claims priority of U.S. provisional patent application Serial No. 60/328,162, filed October 9, 2001, and U.S. provisional patent application Serial No. 60/328,689, filed October 10, 2001, both of which are  
10 incorporated by reference herein.

### BACKGROUND OF THE INVENTION

This invention was made with the assistance of the National Institutes of Health under Grant Nos. GM23200 and CA81534. The U.S. Government has certain rights in this invention.

15 The cytokines IL-4 and IL-13 interact with receptors on target B cells, and stimulate the production of IgE and other mediators of allergy. However, recent data indicate that IL-4/IL-13 signaling also (1) inhibits apoptosis in malignant B cells and other cancer cells, (2) prevents the rejection of tumors by the body, (3) promotes the survival of fibroblasts and therefore increases  
20 fibrosis, and (4) stimulates the differentiation of antigen-presenting cells.

The STAT4 and STAT6 genes encode transcription factors that when phosphorylated by Janus kinases are activated and transported to the nucleus where they regulate cytokine-induced gene expression. See, *e.g.*, J. T. Ihle, Stem Cells Suppl., 1, 105 (1997); M. Heim, J. Recept. Signal. Transduction Res., 19, 75 (1999); K. S. Liu et al., Curr. Opin. Immunol., 10, 271 (1998). For example,  
25 STAT-6 is the common transcription factor for IL-4 and IL-13.

STAT4 and STAT6 are essential for the development of CD4<sup>+</sup> Th1 and Th2 development, respectively. Tumor immunologists have hypothesized that Th1 cells are critical in tumor immunity because they facilitate differentiation of  
30 CD8<sup>+</sup> T cells, which are potent anti-tumor effectors. S. Ostrand-Rosenberg et al., J. Immunol., 165, 6015 (2000) used STAT4<sup>-/-</sup> and STAT6<sup>-/-</sup> mice to test this hypothesis. BALB/c and knockout mice were challenged in the mammary gland with the highly malignant and spontaneously metastatic BALB/c-derived 4T1

mammary carcinoma. Primary tumor growth and metastatic disease were reduced in STAT6<sup>-/-</sup> mice relative to BALB/c and STAT4<sup>-/-</sup> mice. Ab depletions demonstrated that the effect is mediated by CD8<sup>+</sup> T cells, and immunized STAT6<sup>-/-</sup> mice had higher levels of 4T1-specific CTL than BALB/c or STAT4<sup>-/-</sup> mice. Th1 or Th2 cells were not involved, because CD4 depletion did not diminish the anti-tumor effect. Therefore, deletion of the STAT6 gene facilitates development of potent anti-tumor immunity via a CD4<sup>+</sup>-independent pathway.

Sumitomo Pharmaceutical Co. (published Japanese Patent Application, JP 1997/000288026) discloses certain imidazo [2,1-b]thiazole derivatives that are capable of inhibiting STAT-6. The compounds are disclosed to be useful for the treatment and prevention of allergic diseases and parasitic infectious diseases. However, a continuing need exists for small molecules that can inhibit STAT-6 and thus, inhibit IL-4 and IL-13 signal transduction. Such compounds can be used therapeutically as discussed hereinbelow.

In addition, there is a need for novel, potent, and selective agents to prevent detrimental effects upon cells due to DNA damage, such as caused by chemotherapy, radiation, ischemic event, including ischemia-reperfusion injury and organ transplantation, and the like. There is also a need for pharmacological tools for the further study of the physiological processes associated with intracellular DNA damage.


p53, the product of the p53 tumor suppressor gene, is a multifunctional tumor suppressor protein, involved in the negative control of cell growth. In response to a variety of stressors, p53 induces growth arrest or apoptosis, thereby eliminating damaged and potentially dangerous cells. T. M. Gottleib et al., Biochim. Biophys. Acta, 1287, 77 (1996). Mutations in the p53 gene are frequently associated with the metastatic stage of tumor progression, and lack of functional p53 is accompanied by rapid tumor progression, resistance to anti-cancer therapy and increased tumor angiogenesis. See, *e.g.*, A. J. Levine et al., Br. J. Cancer, 69, 409 (1994); R. J. Steele et al., Br. J. Surg., 85, 1460 (1998); C. Cordon-Cardo et al., Surg. Oncol., 13, 319 (1997). p53 deficiency in mice is associated with a high frequency of spontaneous cancers. L. A. Donehower et

al., Nature, 356, 215 (1992); T. Jacks et al., Curr. Biol., 4, 1 (1994). On the basis of these reports, the inactivation of p53 was viewed as an unfavorable event, and it has been speculated that cancer can be inhibited by restoration of p53 function.

- 5 A continuing need exists for compounds that can protect mammalian cells from the damaging effects of chemotherapy and irradiation, or in other situations in which it is desirable to protect tissue from the consequences of clinical or environmental stress.

## 10 SUMMARY OF THE INVENTION

- The present invention provides compounds that act to inhibit the activity of STAT-6 in mammalian cells, and a method to effectively inhibit signal transduction through the IL-4 and IL-13 pathways, *in vitro* or *in vivo*, in the cells of a mammal, such as a human, subject to pathology that is ameliorated by such inhibition. Accordingly, there is provided a method of suppression comprising  
15 administering to a mammal in need of said suppression an effective amount of a compound of formula (I):



(I)


- 20 wherein R<sup>1</sup>, R<sup>2</sup> and R<sup>3</sup> are independently hydrogen, halo, hydroxy, cyano, , -N(R<sub>a</sub>)(R<sub>b</sub>), -S(R<sub>a</sub>), -NO<sub>2</sub>, (C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy, (C<sub>2</sub>-C<sub>6</sub>)alkynyl, (C<sub>2</sub>-C<sub>6</sub>)-alkenyl, (C<sub>2</sub>-C<sub>7</sub>)alkanoyl, (C<sub>2</sub>-C<sub>7</sub>)alkanoyloxy, or (C<sub>3</sub>-C<sub>7</sub>)cycloalkyl or R<sup>1</sup> and R<sup>2</sup> taken together are benzo, optionally substituted by R<sup>1</sup>, or are (C<sub>3</sub>-C<sub>5</sub>)alkylene or methylenedioxy; wherein R<sub>a</sub> and R<sub>b</sub> are each independently hydrogen, (C<sub>1</sub>-  
25 C<sub>3</sub>)alkyl, (C<sub>2</sub>-C<sub>4</sub>)alkanoyl, phenyl, benzyl, or phenethyl; or R<sub>a</sub> and R<sub>b</sub> together with the nitrogen to which they are attached are a 5-6 membered heterocyclic ring, preferably a pyrrolidino, piperidino or morpholino ring;

Ar is aryl, heteroaryl, or a 5-6 membered heterocyclic ring, preferably comprising 1-3 N(R<sub>a</sub>), nonperoxide O or S atoms, such as a pyrrolidino, piperidino or morpholino ring, optionally substituted with 1-5, preferably 1-2, halo, CF<sub>3</sub>, hydroxy, CN, -N(R<sub>a</sub>)(R<sub>b</sub>), (C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy, (C<sub>2</sub>-C<sub>7</sub>)alkanoyl, (C<sub>2</sub>-C<sub>7</sub>)alkanoyloxy, (C<sub>3</sub>-C<sub>7</sub>)cycloalkyl, (C<sub>2</sub>-C<sub>6</sub>)alkenyl, or phenyl groups;

Y is oxy (-O-), -S(O)<sub>0-2</sub>-, Se, -C(R<sup>1</sup>)(R<sup>3</sup>)-, -N(R<sub>a</sub>)-, or -P-; or a pharmaceutically acceptable salt thereof.

Preferably, Ar is not substituted with halo or alkoxy. Preferably, Ar is heteroaryl or a heterocyclic ring. Preferably, R<sup>1</sup> and R<sup>2</sup> are not benzo or (C<sub>3</sub>-C<sub>5</sub>)alkylidenyl when Ar is aryl, *e.g.*, is phenyl or naphthyl. Novel compounds of formula (I) are also within the scope of the present invention, *e.g.*, preferably Y is -O-, -Se-, -C(R<sub>1</sub>)(R<sub>3</sub>)-, or P. Preferably, Ar is heteroaryl. Preferably, Ar is substituted with CN, (C<sub>2</sub>-C<sub>7</sub>)alkanoyl, (C<sub>2</sub>-C<sub>7</sub>)alkanoyloxy, (C<sub>3</sub>-C<sub>7</sub>)cycloalkyl, (C<sub>2</sub>-C<sub>6</sub>)alkenyl or combinations thereof. Preferably, R<sup>1</sup>, R<sup>2</sup> and R<sup>3</sup> are independently, OH, CN, -N(R<sub>a</sub>)(R<sub>b</sub>), -S(R<sub>a</sub>), -NO<sub>2</sub>, (C<sub>2</sub>-C<sub>7</sub>)alkanoyl, or (C<sub>2</sub>-C<sub>7</sub>)alkanoyloxy.

The present method also provides a therapeutic method comprising suppressing STAT-6 or the IL-4/IL-13 pathways in mammalian cells *in vitro* or *in vivo*, and thus treating a pathological condition ameliorated by said suppression, comprising administering to a mammal in need of said suppression an effective amount of a compound of formula (II):




(II)

wherein R<sub>1</sub>, R<sub>2</sub> and R<sub>3</sub> as well as Ar are defined as above; R<sub>4</sub> is the same as, but independent from, R<sub>1</sub>, R<sub>2</sub> and R<sub>3</sub>. R<sub>4</sub> in combination with R<sub>1</sub> can also be benzo,


(C<sub>3</sub>-C<sub>5</sub>)alkylidene or methylenedioxy. These compounds are imidazo[1,2-a]-quinazolines.

Compounds of formula (II) also include (IIa) and (IIb):



- 5 wherein R<sub>1</sub>, R<sub>2</sub>, R<sub>3</sub> and R<sub>4</sub> are as defined herein. Novel compounds of formulae II, IIa and IIb are also within the scope of the invention. Preferably, R<sub>4</sub> is not OH in IIa or IIb, *e.g.*, where R<sub>1</sub> and R<sub>2</sub> or R<sub>1</sub> and R<sub>4</sub> are benzo. In compounds of formula II, R<sub>1</sub> and R<sub>2</sub> are preferably not benzo when Ar is phenyl.

The present invention also includes compounds of formula III:




10

wherein R<sub>1</sub>, R<sub>2</sub> and R<sub>4</sub>, as well as Ar are defined as herein, for formula (I).

- Also included within the invention are methods of using compounds of formula III in amounts effective to suppress STAT-6 or the IL-4/IL-13 pathways  
 15 in mammalian cells, and thus to provide treatment for a mammal afflicted by a pathology ameliorated by said suppression.


Compounds of formula (IV) are also included in the invention:



(IV)

wherein  $R_1$ ,  $R_2$  and  $R_4$ , as well as Ar are defined as above, for formula (II), as well as methods for their use to treat conditions ameliorated by a suppression of  
 5 STAT-6 or by inhibition of signal transduction through the IL-4/IL-13 pathways in mammalian cells *in vitro* or *in vivo*. Preferably,  $R_1$  and  $R_2$  are not benzo when  $R_4$  is H or OH.

Compounds of formula (V) are also included in the invention:



(V)

10

wherein  $R_1$ ,  $R_2$ ,  $R_3$  and  $R_4$  as well as Ar are defined as above, for formula (II), as well as methods for their use as discussed above. Preferably, Ar is not 4-methoxyphenyl when  $R_1$  and  $R_2$  are benzo and  $R_4$  is H.

Compounds of formulae (I)-(V) are small molecule antagonists of IL-  
 15 4/IL-13 signal transduction in mammalian cells *in vitro* and *in vivo*. These molecules can inhibit the survival of malignant B cells and sensitize them to other chemotherapeutic agents, such as, for example, interferons, particularly  $\alpha$ -interferon,  $\beta$ -interferon and  $\gamma$ -interferon. These compounds are relatively nontoxic to normal lymphocytes. Antibodies to IL-4 and IL-13 receptors and to  
 20 other receptors are in clinical trials. However, IL-4 and IL-13 have redundant activities, and thus blocking only one of them is insufficient in many instances. Preferred compounds (I)-(IV) can block both IL-4 and IL-13 signaling. They may act by inhibiting expression of the STAT-6 gene, and thus by inhibiting

STAT-6, the common transcription factor for IL-4 and IL-13. They can be useful to treat cancer, fibrotic diseases and inflammatory diseases.

More specifically, compounds (I)-(V) may be useful for:

1. Treatment of leukemia, lymphoma, Hodgkin's, lung, head, neck,  
5 glioblastomas and other cancers expressing IL-4 and/or IL-13 receptors  
(*e.g.*, gliomas and head and neck cancers).
2. Sensitization of cancer cells to monoclonal antibodies and  
chemotherapeutic agents.
3. Use in vaccines against cancer and viral diseases to increase cytotoxic  
10 T cell responses.
4. Treatment of proliferative fibrotic diseases, such as rheumatoid arthritis,  
pulmonary fibrosis, liver cirrhosis, and chronic kidney diseases.
5. Use alone or in combination with therapeutic agents for treatment of viral  
15 diseases such as hepatitis, papilloma or RNA viruses such as Semliki  
Forest virus, San Angelo Virus, Punta Toro virus, Banzi virus and the  
like.
6. Use alone or in combination with therapeutic agents for treatment of  
autoimmune diseases such as autoimmune diseases, such as lupus  
erythematosus, multiple sclerosis, infertility from endometriosis, type I  
20 diabetes mellitus, Crohn's disease, ulcerative colitis, inflammatory bowel  
disease, and rheumatoid arthritis or diseases such as pulmonary fibrosis.

IL-4 and IL-13 are known to be essential for asthma and allergies. T. Akimoto et al., J. Exp. Med., 182, 1537 (1998) report that STAT-6 deficient mice, which cannot respond to IL-4/IL-13, also do not develop allergic asthma.

25 M. Dancescu et al., J. Exp. Med., 176, 1319 (1992) and U. Kapp, J. Exp. Med., 189, 1939 (1999) report that IL-4 and IL-13 are survival factors for malignant cells in chronic lymphocytic leukemia and Hodgkin's disease (a form of lymphoma). Thus, the present compounds should be useful for treatment of these diseases.

30 K. Kawakami et al., Cancer Res., 60, 2981 (2000) reports the expression of IL-4 receptors in head and neck cancer, melanoma, breast cancer, ovary



cancer, neuroblastomas, renal carcinomas. The present compounds thus can be useful for treatment of these cancers.

M. Terabe et al., Nature/Immunol., 1, 516 (2000) and S. Ostrand-Rosenberg, cited above, report the remarkable finding that lack of STAT-6  
5 signaling promotes immune rejection of cancers. Thus, the claimed compounds can be used in cancer vaccines and/or with monoclonal antibodies to enhance their immunologic effects.

U. Muller-Ladner et al., J. Immunol., 164, 3894 (2000) reported that the  
10 IL-4 pathway is active in the fibroblasts that show unrestrained growth in the joints of patients with rheumatoid arthritis. Similar outgrowth of fibroblasts is seen in pulmonary fibrosis, cirrhosis, renal diseases, scleroderma. The present compounds can be useful in all these conditions.

The invention also provides pharmaceutical compositions comprising novel compounds of formula (I)-(V), or a pharmaceutically acceptable salt  
15 thereof, in combination with a pharmaceutically acceptable diluent or carrier.

The invention also provides novel compounds of formula (I), or a pharmaceutically acceptable salt thereof, in combination with a pharmaceutically acceptable diluent or carrier. Such compounds can be represented by compounds of formula (I), with the proviso that when Y is S, Ar is not phenyl  
20 (C<sub>6</sub>H<sub>5</sub>).

Additionally, the invention provides a therapeutic method for preventing or treating a pathological condition or symptom in a mammal, such as a human, wherein the activity of STAT-6 or IL-4/IL-13-mediated signal transduction is implicated and antagonism or suppression of their action is desired, comprising  
25 administering to a mammal in need of such therapy, an effective amount of one or more compounds of formula (I)-(V), or a pharmaceutically acceptable salt thereof. Such pathological conditions or symptoms include treatment of cancers expressing IL-4 and/or IL-13 receptors, sensitization of cancer cells to chemotherapy or radiation, increasing T<sub>c</sub> cell responses and the treatment of  
30 proliferative fibrotic disease.

The invention provides a compound of formula (I)-(V) for use in medical therapy as well as the use of a compound of formula (I)-(V) for the manufacture

of a medicament for the treatment of a pathological condition or symptom in a mammal, such as a human, which is associated with STAT-6 activation, activation of the IL-4 and/or IL-13 pathways, or p53-induced cellular damage, *i.e.*, with unwanted apoptosis.

5           The invention also includes a method for binding a compound of formula (I)-(V) to cells and biomolecules comprising IL-4 and/or IL-13 receptors, *in vivo* or *in vitro*, comprising contacting said cells or biomolecules with an amount of a compound of formula (I)-(V) effective to bind to said receptors. Cells or biomolecules comprising ligand-bound IL-4/IL-13 receptor sites can be used to  
10   measure the selectivity of test compounds for specific receptor subtypes, or can be used as a tool to identify potential therapeutic agents for the treatment of diseases or conditions associated with IL-4/IL-13 pathway activation, by contacting said agents with said ligand-receptor complexes, and measuring the extent of displacement of the ligand and/or binding of the agent, by methods  
15   known to the art.

          In another embodiment, the present invention provides a compound of formula (I)-(V) that acts to suppress p53 activity in mammalian cells, and a method to effectively suppress p53 activity in the cells of a mammal subject to a stress or pathology that is ameliorated by such suppression. Accordingly, there  
20   is provided a method of p53 suppression comprising administering to a mammal in need of said suppression an effective amount of a compound of formula (I)-(V).

          The invention also provides novel p53 suppressor compounds, as well as pharmaceutical compositions comprising novel compounds of formula (I)-(V),  
25   or a pharmaceutically acceptable salt thereof, in combination with a pharmaceutically acceptable diluent or carrier. Such compounds can be represented by compounds of formula (I), with the proviso that when Y is S, Ar is not phenyl (C<sub>6</sub>H<sub>5</sub>).

          In another embodiment, the present invention provides a compound of  
30   formula (I)-(V) that acts to suppress the IL-4/IL-13 pathway in mammalian cells and a method to effectively suppress the IL-4/IL-13 pathway in the cells of a mammal subject to treatment with cancer agents. This suppression enhances the

efficacy of and thus, could allow the use of lower doses of the cancer agents, which can reduce side effects from the agents.

In another embodiment, the present invention provides a method for the selection of tumors that respond to antagonism of the IL-4/IL-13 pathways using  
5 the compounds of the invention. The method comprises *in vitro* testing of tissue samples to determine if the IL-4/IL-13 pathway is active by measuring the status of STAT-6 in the cell. The presence of phosphorylated STAT-6 is a marker of an active IL-4/IL-13 pathway (see for example, Takeda, K, et al., "The Essential Role of STAT-6 in IL-4 Signalling", *Nature*, 380:627-620, 1996. Thus, the  
10 method specifically comprises measuring phosphorylated STAT-6 in tissue samples from biopsies or isolated cells using phospho-STAT-6 antibodies (*e.g.*, commercially available antibody 575144 from CalBiochem) by immunohistochemistry or immunoblotting.

Additionally, the invention provides a therapeutic method for preventing  
15 or treating a pathological condition or symptom in a mammal, such as a human, wherein the activity of p53 is implicated and antagonism or suppression of its action is desired, comprising administering to a mammal in need of such therapy, an effective amount of a compound of formula (I)-(V), or a pharmaceutically acceptable salt thereof. Such pathological conditions or  
20 symptoms include blocking, moderating or reversing the deleterious effects of chemotherapeutic agents, particularly those which damage DNA; radiation, particularly radiation therapy (gamma-, beta- or UV-radiation), ischemic event, including stroke, infarct, ischemia-reperfusion injury and ischemia due to organ, tissue or cell transplantation; environmental pollution or contamination and the  
25 like.

The invention also includes a method for binding a compound of formula (I) to cells and biomolecules comprising p53 receptors, *in vivo* or *in vitro*, comprising contacting said cells or biomolecules with an amount of a compound of formula (I) effective to bind to said receptors. Cells or biomolecules  
30 comprising ligand-bound p53 receptor sites can be used to measure the selectivity of test compounds for specific receptor subtypes, or can be used as a tool to identify potential therapeutic agents for the treatment of diseases or

conditions associated with p53 activation, by contacting said agents with said ligand-receptor complexes, and measuring the extent of displacement of the ligand and/or binding of the agent, by methods known to the art.

As used herein, the term "p53" or "p53 activity" refers to p53 protein.

- 5 The invention is believed to work by temporarily suppressing expression of the p53 gene and/or activity of p53 protein.

### BRIEF DESCRIPTION OF THE FIGURES

Figure 1 depicts the effects of IBT and PFT- $\alpha$  on B-CLL viability.

- 10 Figure 2 depicts the protective effect of IBT against spontaneous apoptosis and against fludarabine-induced apoptosis.

Figure 3 shows the ability of the various compounds to block the expression of a STAT-6 dependent reporter gene.

- 15 Figure 4 shows the ability of compounds of the invention to reduce the survival of malignant B cells from a patient with chronic lymphocytic leukemia maintained in tissue culture for 72 hours.

Figure 5 shows the structures of compounds numbered in Figures 3-4. Compound 8 is IBT (control).

Figure 6 shows the structures of compounds numbered in Figures 7-11.

- 20 Figure 7 shows the Inhibition of STAT-6 using compound 8, IBT.

Figures 8 and 9 illustrate the Anti-STAT-6 Activity of IBT (compound 8) at several concentrations.

Figure 10 illustrates a functional measurement of Anti-STAT-6 Activity of compounds of the invention.

- 25 Figure 11 illustrates interferon regulation using IBT.

Figure 12 illustrates the effect of IBT and IFN on L1236 Cells

### DETAILED DESCRIPTION

- 30 The following definitions are used, unless otherwise described: halo is fluoro, chloro, bromo, or iodo. Alkyl, alkoxy, alkenyl, alkynyl, etc. denote both straight and branched groups; but reference to an individual radical such as "propyl" embraces only the straight chain radical, a branched chain isomer such

as "isopropyl" being specifically referred to. Aryl denotes a phenyl radical or an ortho-fused bicyclic carbocyclic radical having about nine to ten ring atoms in which at least one ring is aromatic. Heteroaryl encompasses a radical attached via a ring nitrogen or carbon of a monocyclic aromatic ring containing five or six  
5 ring atoms consisting of carbon and one to four heteroatoms each selected from the group consisting of non-peroxide oxygen, sulfur, and N(X) wherein X is absent or is H, O, (C<sub>1</sub>-C<sub>4</sub>)alkyl, phenyl or benzyl. Heteroaryl also includes a radical of an ortho-fused bicyclic heterocycle of about eight to ten ring atoms, particularly a benzo-derivative or one derived by fusing a propylene,  
10 trimethylene, or tetramethylene diradical thereto. Preferred heteroaryls include pyridin-4-yl and thiophen-2-yl. The term "heterocyclic ring" "heterocycle," or "heterocycyl," is defined as above for formula (I).

It will be appreciated by those skilled in the art that compounds of the invention having a chiral center may exist in and be isolated in optically active  
15 and racemic forms. Some compounds may also exhibit polymorphism. It is to be understood that the present invention encompasses any racemic, optically active, polymorphic, or stereoisomeric form, or mixtures thereof, of a compound of the invention, which possess the useful properties described herein, it being well known in the art how to prepare optically active forms (for example, by  
20 resolution of the racemic form by recrystallization techniques, by synthesis from optically active starting materials, by chiral synthesis, or by chromatographic separation using a chiral stationary phase) and how to determine STAT-6 suppression activity using the standard tests described herein, or using other similar tests which are well known in the art. When R<sup>4</sup> is OH, enol or keto  
25 forms of compounds (II)-(V) are also within the scope of the invention, wherein the adjacent N may be replaced by -N(R<sub>a</sub>).

Specific and preferred values listed below for radicals, substituents, and ranges, are for illustration only; they do not exclude other defined values or other values within defined ranges for the radicals and substituents.

30 Specifically, (C<sub>1</sub>-C<sub>6</sub>)alkyl can be methyl, ethyl, propyl, isopropyl, butyl, iso-butyl, sec-butyl, pentyl, 3-pentyl, or hexyl; (C<sub>3</sub>-C<sub>7</sub>)cycloalkyl can be cyclopropyl, cyclobutyl, cyclopentyl, or cyclohexyl; the term cycloalkyl includes

(cycloalkyl)alkyl of the designated number of carbon atoms; (C<sub>1</sub>-C<sub>6</sub>)alkoxy can be methoxy, ethoxy, propoxy, isopropoxy, butoxy, iso-butoxy, sec-butoxy, pentoxy, 3-pentoxy, or hexyloxy; (C<sub>2</sub>-C<sub>6</sub>)alkenyl can be vinyl, allyl, 1-propenyl, 2-propenyl, 1-butenyl, 2-butenyl, 3-butenyl, 1-pentenyl, 2-pentenyl, 3-pentenyl, 4-pentenyl, 1-hexenyl, 2-hexenyl, 3-hexenyl, 4-hexenyl, or 5-hexenyl; (C<sub>2</sub>-C<sub>6</sub>)alkynyl can be ethynyl, 1-propynyl, 2-propynyl, 1-butylnyl, 2-butylnyl, 3-butylnyl, 1-pentylnyl, 2-pentylnyl, 3-pentylnyl, 4-pentylnyl, 1-hexynyl, 2-hexynyl, 3-hexynyl, 4-hexynyl, or 5-hexynyl; (C<sub>2</sub>-C<sub>7</sub>)alkanoyl can be acetyl, propanoyl or butanoyl; (C<sub>2</sub>-C<sub>7</sub>)alkanoyloxy can be acetoxyl, propanoyloxy, butanoyloxy, isobutanoyloxy, pentanoyloxy, or hexanoyloxy; aryl can be phenyl, indenyl, or naphthyl; and heteroaryl can be furyl, imidazolyl, triazolyl, triazinyl, oxazolyl, isoxazolyl, thiazolyl, isothiazolyl, pyrazolyl, pyrrolyl, pyrazinyl, tetrazolyl, pyridyl, (or its N-oxide), thienyl, pyrimidinyl (or its N-oxide), indolyl, isoquinolyl (or its N-oxide) or quinolyl (or its N-oxide).

15 A specific value for R<sup>1</sup> and R<sup>2</sup> is hydroxy, cyano, , -N(R<sub>a</sub>)(R<sub>b</sub>), -S(R<sub>a</sub>), -NO<sub>2</sub>, (C<sub>2</sub>-C<sub>7</sub>)alkanoyl, or (C<sub>2</sub>-C<sub>7</sub>)alkanoyloxy

A specific value for R<sup>1</sup> and R<sup>2</sup> together is butylene or benzo.

A specific value for R<sup>1</sup> and R<sup>4</sup> together is butylene or benzo.

A specific value for R<sup>3</sup> is H.

20 A specific value for R<sup>4</sup> is H.

A specific value for Ar is aryl or heteroaryl, optionally substituted with 1-5, halo, CF<sub>3</sub>, hydroxy, CN, -N(R<sub>a</sub>)(R<sub>b</sub>), (C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy, (C<sub>2</sub>-C<sub>7</sub>)alkanoyl, (C<sub>2</sub>-C<sub>7</sub>)alkanoyloxy, (C<sub>3</sub>-C<sub>7</sub>)cycloalkyl, (C<sub>2</sub>-C<sub>6</sub>)alkenyl, or phenyl groups.

25 Another specific value for Ar is aryl substituted with 1-2, groups.

A specific value for Ar is heteroaryl or phenyl substituted with CN, (C<sub>2</sub>-C<sub>7</sub>)alkanoyl, (C<sub>2</sub>-C<sub>7</sub>)alkanoyloxy, (C<sub>2</sub>-C<sub>7</sub>)cycloalkyl or (C<sub>2</sub>-C<sub>6</sub>)alkenyl.

A specific value for Ar is phenyl, 2, 3 or 4-pyridyl or 2-thienyl; pyrrolidino, piperidino or morpholino.

30 A more specific value for Ar is phenyl, 4-pyridyl or 2-thienyl.

A specific value for Y is oxy (-O-), S(O)<sub>0-2</sub>, C(R<sup>1</sup>)(R<sup>3</sup>), N(R<sub>a</sub>), or -P-.

A specific value for Y is -S-, -O-, -N(R<sub>a</sub>)-, or -P-.

A specific value for Y is -P-, -Se-, -SO-, -SO<sub>2</sub>- or -C(R<sub>1</sub>)(R<sub>3</sub>)-

A specific value for Y is -P-, -Se-, -S(O)- or -SO<sub>2</sub>-.

A more specific value for Y is -S-, -O-, or -NH-,

A specific value for -N(R<sub>a</sub>)(R<sub>b</sub>) is amino.

5 A specific value for -N(R<sub>a</sub>)(R<sub>b</sub>) is pyrrolidino, piperidino or morpholino.

A specific value for halo is Br or F.

A specific method of the invention is treating a disease where the interferon is interferon- $\alpha$ , interferon- $\beta$ , interferon- $\gamma$  or a mixture thereof.

10 Another specific method of the invention is where the interferon is interferon- $\alpha$ .

Another specific method of the invention is where the mammal is a human.

A specific disease for treatment is cancer.

15 A specific cancer for treatment is leukemia, lymphoma, Hodgkin's disease, lung, head, neck, pancreatic or glioblastoma.

A specific leukemia is chronic lymphocytic leukemia (CLL), acute lymphocytic leukemia (ALL), acute myelogenous leukemia (AML), chronic myelogenous leukemia (CML), or multiple myeloma (MM).

20 A more specific leukemia is chronic lymphocytic leukemia, or multiple myeloma.

A specific disease for treatment is for a proliferative fibrotic disease.

A specific proliferative fibrotic disease is rheumatoid arthritis, pulmonary fibrosis, interstitial fibrosis of the lung, scleroderma, keloids, renal fibrosis, liver cirrhosis, endometriosis or chronic kidney disease

25 A specific disease for treatment is for a viral disease.

A specific viral disease is hepatitis, papilloma, semliki forest virus, san angelo virus, punta toro virus, herpes simplex virus, corona virus, cytomegalo virus or Banzi virus.

30 A specific herpes simplex virus is herpes simplex virus type 1 or herpes simplex virus type 2.

A specific disease for treatment is for an autoimmune disease.

A specific autoimmune disease is lupus erythematosus, multiple sclerosis, infertility from endometriosis, type I diabetes mellitus, Crohn's disease, ulcerative colitis, inflammatory bowel disease, rheumatoid arthritis or pulmonary fibrosis.

5 A specific method of the invention is where the method comprises the use of a second therapeutic agent.

Another specific method of the invention is where the therapeutic agent is an antiviral agent, an anticancer agent or a therapeutic agent for treatment of autoimmune diseases.

10 Another specific method of the invention is where the therapeutic agent is ribavirin, acyclovir, valcyclovir or gancyclovir.

A specific method of the invention is administering a mixture of the compounds having formula (I), (II), (III), (IV), (V) or a mixture thereof and interferon delivering a composition to a patient by administering to the patient  
15 the composition of the in.

A specific method of the invention is co-administering the compounds having formula (I), (II), (III), (IV), (V) or a mixture thereof simultaneously with the administration of interferon.


A specific method of the invention is administering the compounds  
20 having formula (I), (II), (III), (IV), or (V) from 0.1 to about 4 hours prior to the administration of interferon.

A specific method of the invention is administering the compounds having formula (I), (II), (III), (IV), (V) or a mixture thereof from 0.1 to about 4 hours after to the administration of interferon.

25 The present invention is based on the discovery that PFT- $\alpha$  is both cytotoxic to mammalian cells and unstable in aqueous solution under *in vivo* conditions. PFT- $\alpha$  undergoes spontaneous ring closure in protic solvents, such as alkanols, to form the imidazo[2,1-b]benzothiazole derivative, abbreviated IBT, as shown in Scheme 1.

30






Scheme 1


Biological evaluation, described below, demonstrated that IBT is actually responsible for the observed p53 inhibition observed by Komarov et al. (*Science*, 285, 1733 (1999)). Thus, since IBT and compounds of formula (I) are expected to be both less toxic and more stable than imino compounds such as PFT- $\alpha$ , they are desirable agents for protection of mammalian cells against a wide variety of stressors, including therapeutic agents, and clinical and environmental trauma.

Compounds of formula (I) can be readily prepared as disclosed by Singh et al., *Indian J. Chem.*, 7, 997 (1996), as shown in Scheme 2.




Scheme 2

In Scheme 2, a suitable 2-aminobenzothiazole derivative is reacted with an  $\alpha$ -haloketone in refluxing ethanol resulting in alkylation and ring closure in one single step. An example for the pyridinyl-substituted derivative is given below:



In Scheme 2, the reaction of 1 and 4 can be carried out simply by combining the compounds in a suitable aprotic solvent such as benzene. See, I. Soldabols et al., Khim. Pharm. Zh., 1, 17 (1967). The conversion of 1 → 3 can also be accomplished in one step by refluxing 1 and the phenacyl bromide 4 in ethanol.

- 5 Singh et al. used starting materials wherein R<sup>1</sup> and R<sup>2</sup> together are -(CH<sub>2</sub>)<sub>4</sub>- or -CH(CH<sub>3</sub>)-(CH<sub>2</sub>)<sub>3</sub>- and Ar is substituted phenyl. Recently, Sumitomo Pharmaceutical Co. Ltd. (Japanese Pat. No. 11-29475) (1999) disclosed the preparation of certain compounds of formula 2, wherein R<sup>3</sup> is H and Ar is substituted phenyl, and Japanese Pat. No. 11-106340 (1999) disclosed the
- 10 preparation of certain compounds of formula 3 wherein Ar is substituted phenyl or naphthyl and R<sup>1</sup> and R<sup>2</sup> can be, *inter alia*, H, alkylene or benzo. Compounds of formula 1 were prepared according to Scheme 3.




Scheme 3

- 15 The compounds of formula (I) are disclosed to be useful for "the treatment and prevention of allergic disease and parasitic infectious diseases, or the like."

Certain of the compounds of formula (I) are useful as intermediates to prepare other compounds of formula (I), as would be recognized by the art.


- 20 Compounds of formulae (II)-(V) can be prepared as generally described in PCT/WO97/42192; U.S. Pat. No. 4,020,062, Armianianskii Khim. Zhuv., 43, 245 (1990); Coppola et al., J. Org. Chem., 41, 825 (1976) (II); M. A. Likhale et al., J. Ind. Chem. Soc., 69, 667 (1992); K. T. Potts et al., J. Org. Chem., 35, 3448 (1970); J. E. Francis et al., J. Med. Chem., 34, 281, 2899 (1991) (IV) and A. Guieffier, J. Het. Chem., 27, 421 (1990) (V).

A general method for preparation of imidazo[1,2-a]quinazolines of formula (II) is found in Coppola, et al., wherein a functionalized isatoic anhydride is first alkylated with the alpha-haloketone and then condensed with a suitable thiopseudourea, as shown below for a pyridinyl derivative:



5


A procedure reported by R. Heckendorn et al., *Helv. Chim. Acta*, **63**, 1 (1980) can be used to prepare the 2-aryl-substituted 1,2,4-triazolo[1,5-a]quinazolines wherein a 2-hydrazinobenzoic acid is condensed with an appropriate N-cyanoimide ester as shown below:



10


A suitable procedure by Francis, et al., cited above, is used to obtain aryl substituted 1,2,4-triazolo[1,5-c]quinazolines of formula (IV), wherein an appropriate anthranilonitrile is converted to the corresponding carbamate by reaction of the nitrile with ethyl carbonate in the presence of sodium ethoxide,

followed by condensation with a suitable aryl carbohydrazide or heteroaryl carbohydrazide as shown below:



5

Imidazo[1,2-c]quinazolines of formula (V) may be prepared according to the procedure outlined by Gueffier, et al., wherein a 4-aminoquinazoline is reacted with a bromomethyl aryl ketone in refluxing ethanol. Heteroaryl ketones may also be used as shown below for a pyridinyl derivative:



10

In cases where compounds are sufficiently basic or acidic to form stable nontoxic acid or base salts, administration of the compounds as salts may be appropriate. Examples of pharmaceutically acceptable salts are organic acid addition salts formed with acids which form a physiological acceptable anion, for example, tosylate, methanesulfonate, acetate, citrate, malonate, tartarate, succinate, benzoate, ascorbate,  $\alpha$ -ketoglutarate, and  $\alpha$ -glycerophosphate.

15

Suitable inorganic salts may also be formed, including hydrochloride, sulfate, nitrate, bicarbonate, and carbonate salts.

Pharmaceutically acceptable salts may be obtained using standard procedures well known in the art, for example, by reacting a sufficiently basic compound such as an amine with a suitable acid affording a physiologically acceptable anion. Alkali metal (for example, sodium, potassium or lithium) or alkaline earth metal (for example, calcium) salts of carboxylic acids can also be made.

The compounds of formula (I)-(V) can be formulated as pharmaceutical compositions and administered to a mammalian host, such as a human cancer patient, in a variety of forms adapted to the chosen route of administration, *i.e.*, orally or parenterally, by intravenous, intramuscular, topical or subcutaneous routes.

Thus, the present compounds may be systemically administered, *e.g.*, orally, in combination with a pharmaceutically acceptable vehicle such as an inert diluent or an assimilable edible carrier. They may be enclosed in hard or soft shell gelatin capsules, may be compressed into tablets, or may be incorporated directly with the food of the patient's diet. For oral therapeutic administration, the active compound may be combined with one or more excipients and used in the form of ingestible tablets, buccal tablets, troches, capsules, elixirs, suspensions, syrups, wafers, and the like. Such compositions and preparations should contain at least 0.1% of active compound. The percentage of the compositions and preparations may, of course, be varied and may conveniently be between about 2 to about 60% of the weight of a given unit dosage form. The amount of active compound in such therapeutically useful compositions is such that an effective dosage level will be obtained.

The tablets, troches, pills, capsules, and the like may also contain the following: binders such as gum tragacanth, acacia, corn starch or gelatin; excipients such as dicalcium phosphate; a disintegrating agent such as corn starch, potato starch, alginic acid and the like; a lubricant such as magnesium stearate; and a sweetening agent such as sucrose, fructose, lactose or aspartame or a flavoring agent such as peppermint, oil of wintergreen, or cherry flavoring

may be added. When the unit dosage form is a capsule, it may contain, in addition to materials of the above type, a liquid carrier, such as a vegetable oil or a polyethylene glycol. Various other materials may be present as coatings or to otherwise modify the physical form of the solid unit dosage form. For instance, 5 tablets, pills, or capsules may be coated with gelatin, wax, shellac or sugar and the like. A syrup or elixir may contain the active compound, sucrose or fructose as a sweetening agent, methyl and propylparabens as preservatives, a dye and flavoring such as cherry or orange flavor. Of course, any material used in preparing any unit dosage form should be pharmaceutically acceptable and 10 substantially non-toxic in the amounts employed. In addition, the active compound may be incorporated into sustained-release preparations and devices.

The active compound may also be administered intravenously or intraperitoneally by infusion or injection. Solutions of the active compound or its salts can be prepared in water, optionally mixed with a nontoxic surfactant. 15 Dispersions can also be prepared in glycerol, liquid polyethylene glycols, triacetin, and mixtures thereof and in oils. Under ordinary conditions of storage and use, these preparations contain a preservative to prevent the growth of microorganisms.

The pharmaceutical dosage forms suitable for injection or infusion can 20 include sterile aqueous solutions or dispersions or sterile powders comprising the active ingredient, which are adapted for the extemporaneous preparation of sterile injectable or infusible solutions or dispersions, optionally encapsulated in liposomes. In all cases, the ultimate dosage form should be sterile, fluid and stable under the conditions of manufacture and storage. The liquid carrier or 25 vehicle can be a solvent or liquid dispersion medium comprising, for example, water, ethanol, a polyol (for example, glycerol, propylene glycol, liquid polyethylene glycols, and the like), vegetable oils; nontoxic glycerol esters, and suitable mixtures thereof. The proper fluidity can be maintained, for example, by the formation of liposomes, by the maintenance of the required particle size 30 in the case of dispersions or by the use of surfactants. The prevention of the action of microorganisms can be brought about by various antibacterial and antifungal agents, for example, parabens, chlorobutanol, phenol, sorbic acid,

thimerosal, and the like. In many cases, it will be preferable to include isotonic agents, for example, sugars, buffers or sodium chloride. Prolonged absorption of the injectable compositions can be brought about by the use in the compositions of agents delaying absorption, for example, aluminum monostearate and  
5 gelation.

Sterile injectable solutions are prepared by incorporating the active compound in the required amount in the appropriate solvent with various of the other ingredients enumerated above, as required, followed by filter sterilization. In the case of sterile powders for the preparation of sterile injectable solutions,  
10 the preferred methods of preparation are vacuum drying and the freeze drying techniques, which yield a powder of the active ingredient plus any additional desired ingredient present in the previously sterile-filtered solutions.

For topical administration, the present compounds may be applied in pure form, *i.e.*, when they are liquids. However, it will generally be desirable to  
15 administer them to the skin as compositions or formulations, in combination with a dermatologically acceptable carrier, which may be a solid or a liquid.

Useful solid carriers include finely divided solids such as talc, clay, microcrystalline cellulose, silica, alumina and the like. Useful liquid carriers include water, alcohols or glycols or water-alcohol/glycol blends, in which the  
20 present compounds can be dissolved or dispersed at effective levels, optionally with the aid of non-toxic surfactants. Adjuvants such as fragrances and additional antimicrobial agents can be added to optimize the properties for a given use. The resultant liquid compositions can be applied from absorbent pads, used to impregnate bandages and other dressings, or sprayed onto the  
25 affected area using pump-type or aerosol sprayers.

Thickeners such as synthetic polymers, fatty acids, fatty acid salts and esters, fatty alcohols, modified celluloses or modified mineral materials can also be employed with liquid carriers to form spreadable pastes, gels, ointments, soaps, and the like, for application directly to the skin of the user.

30 Examples of useful dermatological compositions which can be used to deliver the compounds of formula (I)-(V) to the skin are known to the art; for example, see Jacquet et al. (U.S. Pat. No. 4,608,392), Geria (U.S. Pat. No.

4,992,478), Smith et al. (U.S. Pat. No. 4,559,157) and Wortzman (U.S. Pat. No. 4,820,508).

Useful dosages of the compounds of formula (I)-(V) can be determined by comparing their *in vitro* activity, and *in vivo* activity in animal models.

- 5 Methods for the extrapolation of effective dosages in mice, and other animals, to humans are known to the art; for example, see U.S. Pat. No. 4,938,949.

Generally, the concentration of the compound(s) of formula (I)-(V) in a liquid composition, such as a lotion, will be from about 0.1-25 wt %, preferably from about 0.5-10 wt %. The concentration in a semi-solid or solid composition  
10 such as a gel or a powder will be about 0.1-5 wt %, preferably about 0.5-2.5 wt %.

The amount of the compound, or an active salt or derivative thereof, required for use in treatment will vary not only with the particular salt selected but also with the route of administration, the nature of the condition being  
15 treated and the age and condition of the patient and will be ultimately at the discretion of the attendant physician or clinician.

In general, however, a suitable dose will be in the range of from about 0.5 to about 100 mg/kg, *e.g.*, from about 10 to about 75 mg/kg of body weight per day, such as 3 to about 50 mg per kilogram body weight of the recipient per  
20 day, preferably in the range of 6 to 90 mg/kg/day, most preferably in the range of 15 to 60 mg/kg/day.

The compound is conveniently administered in unit dosage form, for example, containing 5 to 1000 mg, conveniently 10 to 750 mg, most conveniently, 50 to 500 mg of active ingredient per unit dosage form.

25 Ideally, the active ingredient should be administered to achieve peak plasma concentrations of the active compound of from about 0.5 to about 75  $\mu\text{M}$ , preferably, about 1 to 50  $\mu\text{M}$ , most preferably, about 2 to about 30  $\mu\text{M}$ . This may be achieved, for example, by the intravenous injection of a 0.05 to 5% solution of the active ingredient, optionally in saline, or orally administered as a  
30 bolus containing about 1-100 mg of the active ingredient. Desirable blood levels may be maintained by continuous infusion to provide about 0.01-5.0 mg/kg/hr or



by intermittent infusions containing about 0.4-15 mg/kg of the active ingredient(s).


The desired dose may conveniently be presented in a single dose or as divided doses administered at appropriate intervals, for example, as two, three, four or more sub-doses per day. The sub-dose itself may be further divided, *e.g.*, into a number of discrete loosely spaced administrations, such as multiple inhalations from an insufflator or by application of a plurality of drops into the eye.

The ability of a compound of the invention to act as a suppressor of p53 activity may be determined using pharmacological models which are well known to the art, *e.g.*, as disclosed below.

The invention will now be illustrated by the following non-limiting Examples.

### EXAMPLE 1

#### A. Ring-closure of PFT- $\alpha$



The preparation of PFT- $\alpha$  was accomplished as shown in Scheme 1 by reacting 4-methyl-2-bromoacetophenone with 2-amino-4,5,6,7-tetrahydrobenzothiazole. Upon recrystallization of the PFT- $\alpha$  from isopropyl alcohol, it was noticed that PFT- $\alpha$  readily ring-closed completely to the imidazo[2,1-b]benzothiazole (IBT). Therefore, a subsequent investigation was undertaken to study the propensity of PFT- $\alpha$  to ring-close in protic solvents. Initial results indicated that PFT- $\alpha$  begins cyclizing at room temperature immediately upon dissolution in protic solvents. Thus, PFT- $\alpha$  was dissolved in DMSO and water

dilutions were made from this stock. Reversed phase HPLC analysis of the solution at 25°C over time gave results as shown in Table 1.

Table 1

<u>Time (h)</u>	<u>% cyclized to IBT</u>
0	5
12	47
24	69
48	92

5

In addition, NMR studies were used to confirm the structure of the known IBT and a time course in DMSO-d<sub>6</sub> also showed spontaneous conversion of PFT- $\alpha$  to IBT, as judged by the appearance of a new aromatic proton signal at  $\delta$  8.50 ppm in the proton spectrum corresponding to the C<sub>3</sub>H proton.

10

B. 2-(Pyridin-4-yl)imidazo[2,1-b]benzothiazole

A mixture of 2-aminobenzothiazole (0.01 mol) and 4-bromoacetylpyridine (0.01 mol) in anhydrous ethanol (100 mL) is refluxed for 5 hours. The reaction mixture is evaporated to dryness in vacuo and the residue is slurried in ice water. The resulting solid is filtered and dried to provide the title compound as the HBr salt in 60% yield.

15

C. 2-(Pyridin-4-yl)imidazo[1,2-a]quinazolin-9-one

Isatoic anhydride (0.01 mol) is treated with sodium hydride (0.012 mol) in dry dimethylacetamide (50 mL) at room temperature for 20 min. and then 4-bromoacetyl-pyridine (0.01 mol) is added and the mixture is stirred at 80 °C for 20 hours. The mixture is cooled and poured into cold, aqueous sodium carbonate (500 mL, saturated) and extracted with ethyl acetate (3 x 200 mL). The organic layer is dried over magnesium sulfate and evaporated to yield the crude alkylated isatoic anhydride which is used directly without further purification for the ring closure procedure. Thus, this ketone intermediate is suspended in acetonitrile (100 mL) containing methyl-2-thiopseudourea (0.012 mol) and sodium carbonate (0.012 mol) and the mixture is refluxed for 30 min. The solvent is then removed in vacuo and replaced with dichloromethane (100 mL). The

20

25

insoluble salts are filtered off and washed with additional solvent, and the filtrate is evaporated to dryness and diglyme (50 mL) is added to the residue. After addition of one pellet of sodium hydroxide to catalyze the reaction, the mixture is refluxed for 2 hours. Upon cooling, a precipitate forms which is filtered,  
5 washed with a small amount of ethyl acetate and recrystallized from methanol or dichloromethane to yield the title compound.

D. 2-(p-Methylphenyl)[1,2,4]-triazolo[1,5-a]quinazolin-5-4H-one

To a cooled solution (0 °C) of N-cyanoarylethylimidate in absolute alcohol (75 mmol in 100 mL EtOH) is added dropwise triethylamine (225 mmol)  
10 over 30 min. and then 75 mmol of 2-hydrazinobenzoic acid hydrochloride is added portionwise keeping the temperature below 3 °C. The mixture is then allowed to warm slowly to room temperature and is stirred overnight. The resulting mixture is cooled and neutralized with conc. HCl and warmed for 3 hours at 80 °C with stirring. The reaction mixture is diluted with water and  
15 cooled to 5 °C. The resulting solid product which separates is filtered off, washed with cold water, then ether and dried to yield the title compound.

E. 2-(Pyridin-4-yl)imidazo[1,2-c]quinazoline

A mixture of 4-aminoquinazoline (0.01 mol) and 4-bromoacetylpyridine (0.01 mol) in anhydrous ethanol (100 mL) is refluxed for 5 hours. The reaction  
20 mixture is evaporated to dryness in vacuo and the residue is slurried in ice water. The resulting solid is filtered and dried to provide the title compound as the HBr salt.

F. 2-(Pyridin-4-yl)1,2,4-triazolo[1,5-c]quinazolin-5(6H)-one

A mixture of the carbamate of anthranilonitrile (prepared by reacting  
25 anthranilonitrile (0.21 mol) with ethyl carbonate (250 mL) in absolute ethanol (500 mL) containing sodium ethoxide, 1.67 mol) is reacted with 4-pyridinecarbohydrazide (one to one equivalence, 55 mmol each) in 2-ethoxyethanol (185 mL) containing tri-n-propylamine (7.4 mL) by heating at reflux for 16 h, cooling, and treating with water gradually to promote  
30 crystallization. After overnight refrigeration, the solid product is collected and recrystallized from ethanol.

## EXAMPLE 2

### Effect of the p53 Inhibitory Compounds on B-CLL Viability

The malignant lymphocytes from two patients with chronic lymphocytic leukemia [CLL] were isolated by ficoll-hypaque sedimentation and suspended at a density of 1 million cells per milliliter in RPMI 1640 medium supplemented with 10% fetal bovine serum. Two hundred microliter aliquots of cells were dispersed in the wells of culture plates containing the indicated final concentrations of either PFT- $\alpha$  ("PFT-open") or IBT (PFT-closed). After 3 days culture, viable cells were enumerated by fluorescence-activated cell sorting [FACS] after staining with propidium iodide [PI]. Viable cells excluded the dye [open circles]. In addition, cell metabolism was assessed by the ability of the cells to exclude the tetrazolium dye MTT [closed squares]. As shown in Figure 1, the PFT-open dose-dependently reduced CLL survival, whereas PFT-closed [*i.e.*, IBT] was non-toxic at concentrations up to 100 micromolar.

15

## EXAMPLE 3

### Protection Against Spontaneous Apoptosis and

### Apoptosis Induced by the Anti-metabolite Fludarabine

Chronic lymphocytic leukemia [CLL] cells were cultured for 3 days as described in Example 2. Some of the cultures were supplemented with one micromolar of PFT-open or PFT-closed, as indicated. In the experiment shown in the bottom panel of Figure 2, some of the cultures also contained the cytotoxic adenine nucleoside analog fludarabine [abbreviated F-AraA]. Fludarabine is the first line treatment for CLL, and the toxicity of the drug is dependent upon the p53 pathway. To assess healthy, viable cells, staining was done with both PI, as indicated in Example 2, and with the mitochondrial dye DiOC6. Cells that were both PI negative and DiOC6 high were enumerated by FACS. While PFT- $\alpha$  and IBT exhibited nearly equivalent effects on untreated CLL cells, IBT exerted less protective effects when combined with CLL cells treated with F-AraA than did PFT- $\alpha$ .

30

**EXAMPLE 4****Screening of compounds of formula (I) for inhibition of IL-4 transcriptional activity**

5 The BEAS-2B human airway epithelial cells were transiently transfected with the human 12/15-lipoxygenase promoter/luciferase reporter gene. Cells were then incubated with the IBT analogs (Fig. 5) at 10  $\mu$ M for 1 hour, followed by IL-4 (10 ng/ml). After 16 hours, luciferase was measured using a chemiluminometer. The STAT-6 induction was normalized using the B-gal results as "background." The viability of the treated cells was visually verified  
10 at the end of the incubation, and found to be >95%. Results shown in Figure 3 are the mean of duplicate measurements.

**EXAMPLE 5****Sensitization of CLL cells to apoptosis by IL-4/IL-13 antagonists**

15 Chronic lymphocytic leukemia (CLL) cells were isolated from whole blood of patients, cultured in RPMI-1640 supplemented with 10% FB. CLL cells were pre-incubated for 1 hour with the indicated analogs (Fig. 5) at 1  $\mu$ M and exposed for 24 hours to the nucleoside analogs Fludarabine (Fludara) and Cladribine (2 CdA) at 1 and 10  $\mu$ M. Cells were then incubated for 10 minutes in  
20 growing medium with 5  $\mu$ g/ml Propidium iodide and 40 nM DiOC<sub>6</sub> and analyzed by flow cytometry. Viable cells (Y axis) and high DiOC<sub>6</sub> (FL-1) and low PI (FL-3) fluorescence.

**EXAMPLE 6****Preparation of Pharmaceutical Dosage Forms**

25 The following illustrate representative pharmaceutical dosage forms, containing a compound of formula (I)-(V), for therapeutic or prophylactic use in humans.

<u>(i) Table 1</u>	<u>mg/tablet</u>
Compound of Formula (I)-(V)	100.0
Lactose	77.5
Povidone	15.0
Croscarmellose sodium	12.0
Microcrystalline cellulose	92.5
Magnesium stearate	<u>3.0</u>
	300.0

<u>(ii) Table 2</u>	<u>mg/tablet</u>
Compound of Formula (I)-(V)	20.0
Microcrystalline cellulose	410.0
Starch	50.0
Sodium starch glycolate	15.0
Magnesium stearate	<u>5.0</u>
	500.0

<u>(iii) Capsule</u>	<u>mg/capsule</u>
Compound of Formula (I)-(V)	10.0
Colloidal silicon dioxide	1.5
Lactose	465.5
Pregelatinized starch	120.0
Magnesium stearate	<u>3.0</u>
	600.0

<u>(iv) Injection 1 (1 mg/ml)</u>	<u>mg/ml</u>
Compound of Formula (I)-(V)	1.0
Dibasic sodium phosphate	12.0
Monobasic sodium phosphate	0.7
Sodium chloride	4.5
01 N Sodium hydroxide solution	
(pH adjustment to 7.0-7.5)	q.s.

Water for injection	q.s. ad 1 mL
---------------------	--------------

<u>(v) Injection 2 (10 mg/ml)</u>	<u>mg/ml</u>
Compound of Formula (I)-(V)	10.0
Monobasic sodium phosphate	0.3
Dibasic sodium phosphate	1.1
Polyethylene glycol 400	200.0
01 N Sodium hydroxide solution	
(pH adjustment to 7.0-7.5)	q.s.
Water for injection	q.s. ad 1 mL

<u>(vi) Aerosol</u>	<u>mg/can</u>
Compound of Formula (I)-(V)	20.0
Oleic acid	10.0
Trichloromonofluoromethane	5,000.0
Dichlorodifluoromethane	10,000.0
Dichlorotetrafluoroethane	5,000.0

The above formulations may be obtained by conventional procedures well known in the pharmaceutical art.

5

## EXAMPLE 7

### Inhibition of STAT-6

Human airway epithelial cells, BEAS-2B, were transiently transfected with the human 12/15-lipoxygenase promoter/luciferase reporter plasmid with and without a STAT6 expressing plasmid. The cells were incubated with 10  $\mu$ M of STAT-6 (Control Sample 1); 10  $\mu$ M STAT-6 (Sample 2); a mixture of STAT-6 and compound 8 (Sample 3); and compound 8 alone (Sample 4); for 1 hour. This was followed treatment with 10 ng/mL of IL-4 (except control). After 16 hours, luciferase was measured using a chemiluminometer. The STAT-6 induction was normalized using the  $\beta$ -gal results as "background". The viability

of the treated cells was visually verified at the end of the incubation, and found to be >95%. The results are illustrated in Figure 7.

#### EXAMPLE 8

##### 5        Measurement of Anti-STAT-6 Activity of IBT (compound 8)

Human airway epithelial cells, BEAS-2B, were transiently transfected with the human 12/15-lipoxygenase promoter/luciferase reporter gene. The cells were incubated with compound 8 at concentrations of 0.075  $\mu$ M, 0.5  $\mu$ M, 1.0  $\mu$ M, 7.5  $\mu$ M and 40  $\mu$ M for 1 hour. The treatment was followed by incubation  
10 with IL-4 (10 ng/mL). After 16 hours, luciferase was measured using a chemiluminometer. The STAT-6 induction was normalized using the  $\beta$ -gal results as "background". The results are the mean of triplicate measurements, all standard deviations were lower than 8%. The results are illustrated in Figure 8.

15

#### EXAMPLE 9

##### Measurement of Anti-STAT-6 Activity

The inhibitory activity of the compounds of the invention was tested at a fixed concentration of 10  $\mu$ M in IL-4 stimulated BEAS-2B human airway cells using the 12/15-lipoxygenase promoter/luciferase as described in the previous  
20 Examples. The results are illustrated in Figure 9.

#### EXAMPLE 10

##### Functional Measurement of Anti-STAT-6 Activity

The effectiveness of compounds 8, 12 and 28 on the IL-4-induced expression of the protein 12/15 lipooxygenase (12/15-LOX) was measured in  
25 primary human monocytes. The cells were pre-incubated for 1 hour with 10  $\mu$ M of the test compound. This was followed by exposure to 10 ng/ml of IL-4. The expression of 12/15-LOX was quantified by immunoblotting using a specific monoclonal antibody and compared to a loading control. The results are illustrated in Figure 10.

30

#### EXAMPLE 11

##### Interferon Regulation Using IBT (Compound 8)



The B cells RAMOS were incubated with IL-4 (Fig 11B); leukocyte-purified interferons (IFN,  $10^4$  U/ml) (Fig 11C); INF AND IL-4 (Fig 11D); compound 8 alone (Fig 11E); and leukocyte-purified interferons (IFN,  $10^4$  U/ml) (Fig 11E); IL-4 (10 ng/ml) and compound #8 (10  $\mu$ M) (Fig 11F); for 48 hrs. The control is illustrated in Fig 11A. The surface expression of the Fas receptor (also known as FasR or CD95) was then measured by flow cytometry using a PE-conjugated monoclonal antibody. The results as a percentage of CD95+ cells are indicated in each panel in Figure 11.

### EXAMPLE 12

#### Effect of IBT, and Interferon on Hodgkin's disease.

Human Hodgkin's lymphoma cells L1236 were incubated for 3, 5 or 7 days in complete RPMI1640 and 10% FBS. The cells were plated at  $3 \times 10^5$  cells/mL. These cells (L1236) are similarly to other Hodgkin's cell lines and other tumor types such as pancreatic cancers, which express high levels of IL-4 and/or IL-13 receptors or have been shown to require IL-13 for their survival.

The L1236 cells were cultured in presence of type 1 interferon (leukocyte interferon, primarily IFN- $\alpha$ ), or in presence of IBT alone at doses up to 10  $\mu$ M, only a small anti-proliferative effect was observed (Figure 12, second and third columns). When the L1236 cells were cultured in presence of interferon and IBT combined (Figure 12, first column), a stronger growth-inhibitory activity was observed. The effect was more pronounced after 7 days of incubation. These results indicate that IBT or related molecules by inhibiting that the IL4 and/or IL-13 signaling can enhance the effects of interferons. These results suggest that IBT or related analogs will be clinically useful in combination with interferons in diseases where interferons have been shown to be useful such as, but not limited to, cancers and viral infections.


The results are illustrated in Figure 12. The y-axis represent the normalized viability when compared to control cells incubated with 0.1% DMSO with no interferon or IBT added, adjusted at 100%. The cell viability was determined using CellTiter 96 Aqueous One Solution Cell Proliferation Assay (Promega). (CellTiter 96 uses a novel tetrazolium compound (MTS) and an electron coupling reagent, phenazine ethosulfate.)

All publications, patents, and patent documents are incorporated by reference herein, as though individually incorporated by reference. The invention has been described with reference to various specific and preferred  
5 embodiments and techniques. However, it should be understood that many variations and modifications may be made while remaining within the spirit and scope of the invention.

What is claimed is

1. A therapeutic method to enhance the efficacy of interferon treatment comprising administering to a mammal subject to interferon treatment a compound which is an antagonist of the IL-4 or IL-13 signal transduction pathway in an amount effective to enhance said efficacy.
2. The method of claim 1, wherein the interferon is interferon- $\alpha$ , interferon- $\beta$ , interferon- $\gamma$  or a mixture thereof.
3. The method of claim 1 or 2, wherein the interferon is interferon- $\alpha$ .
4. The method of any of claims 1-3, wherein the mammal is a human.
5. The method of any of claims 1-4, wherein the interferon treatment is for cancer.
6. The method of claim 5, wherein the cancer is leukemia, lymphoma, Hodgkin's disease, lung, head, neck, pancreatic or glioblastoma.
7. The method of claim 6, wherein the leukemia is cronic lymphocytic leukemia, acute lymphocytic leukemia, acute myelogenous leukemia, cronic myelogenous leukemia, or multiple myleoma.
8. The method of any of claims 1-4, wherein the interferon treatment is for a proliferative fibrotic disease.
9. The method of claim 8, wherein the proliferative fibrotic disease is rheumatoid arthritis, pulmonary fibrosis, interstitial fibrosis of the lung, scleroderma, keloids, renal fibrosis, liver cirrhosis, or chronic kidney disease.
10. The method of any of claims 1-4, wherein the interferon treatment is for a viral disease.

11. The method of claim 10, wherein the viral disease is hepatitis, papilloma, semliki forest virus, san angelo virus, punta toro virus, herpes simplex virus, corona virus, cytomegalo virus or banzi virus.
12. The method of any of claims 1-4, wherein the interferon treatment is for an autoimmune disease.
13. The method of claim 12, wherein the autoimmune disease is lupus erythematosus, multiple sclerosis, infertility from endometriosis, type I diabetes mellitus, Crohn's disease, ulcerative colitis, inflammatory bowel disease, rheumatoid arthritis or pulmonary fibrosis.
14. The method of any of claims 1-13, further comprising the use of a second therapeutic agent.
15. The method of claim 14, wherein the second therapeutic agent is an antiviral agent, an anticancer agent or a therapeutic agent for treatment of autoimmune diseases.
16. The method of claim 15, where the therapeutic agent is an antiviral agent selected from the group consisting of ribavirin, acyclovir, valcyclovir and gancyclovir.
17. The method of any of claims 1-4, wherein the antagonist of the IL-4 or IL-13 signal transduction pathway is a compound of formula (I):



(I)

wherein R<sup>1</sup>, R<sup>2</sup> and R<sup>3</sup> are independently hydrogen, halo, hydroxy, cyano, -N(R<sub>a</sub>)(R<sub>b</sub>), -S(R<sub>a</sub>), -NO<sub>2</sub>, (C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy, (C<sub>2</sub>-C<sub>6</sub>)alkynyl, (C<sub>2</sub>-C<sub>6</sub>)alkenyl, (C<sub>2</sub>-C<sub>7</sub>)alkanoyl, (C<sub>2</sub>-C<sub>7</sub>)alkanoyloxy, or


(C<sub>3</sub>-C<sub>7</sub>)cycloalkyl or R<sup>1</sup> and R<sup>2</sup> taken together are benzo, optionally substituted by R<sup>1</sup>, (C<sub>3</sub>-C<sub>5</sub>)alkylene or methylene dioxy; wherein R<sub>a</sub> and R<sub>b</sub> are each independently hydrogen, (C<sub>1</sub>-C<sub>3</sub>)alkyl, (C<sub>2</sub>-C<sub>4</sub>)alkanoyl, phenyl, benzyl, or phenethyl; or R<sub>a</sub> and R<sub>b</sub> together with the nitrogen to which they are attached are a 5-6 membered heterocyclic ring;

Ar is aryl, heteroaryl, or a 5-6 membered heterocyclic ring, optionally comprising 1-3 N(R<sub>a</sub>), nonperoxide O or S atoms, optionally substituted with 1-5 groups where the groups are selected from halo, CF<sub>3</sub>, hydroxy, CN, -N(R<sub>a</sub>)(R<sub>b</sub>), (C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy, (C<sub>2</sub>-C<sub>7</sub>)alkanoyl, (C<sub>2</sub>-C<sub>7</sub>)alkanoyloxy, (C<sub>3</sub>-C<sub>7</sub>)cycloalkyl, (C<sub>2</sub>-C<sub>6</sub>)alkenyl, and phenyl;

Y is oxy (-O-), -S(O)<sub>0-2</sub>-, Se, -C(R<sup>1</sup>)(R<sup>3</sup>)-, -N(R<sub>a</sub>)-, or -P-,  
or a pharmaceutically acceptable salt thereof.

18. The method of claim 17, wherein the heterocyclic ring, is pyrrolidino, piperidino or morpholino.
19. The method of claim 17 or 18, wherein the aryl, heteroaryl or heterocyclic group is substituted with 1 or 2 groups.
20. The method any of claims 17-19, wherein R<sup>1</sup> and R<sup>2</sup> together is butylene or benzo.
21. The method of any of claims 17-20, wherein R<sup>3</sup> is hydrogen.
22. The method of any of claims 17-21, wherein Ar is phenyl, 4-pyridyl or 2-thienyl.
23. The method of any of claims 17-22, wherein Ar is a 5-6 membered heterocyclic ring, comprising 1-3 N(R<sub>a</sub>), nonperoxide O or S atoms.
24. The method of any of claims 17-23, wherein Ar is pyrrolidino, piperidino or morpholino.
25. The method of any of claims 17-24, wherein Y is -O- (oxy), -S(O)<sub>0-2</sub>-, -C(R<sup>1</sup>)(R<sup>3</sup>)-, -NR<sup>1</sup>, or -P-.

26. The method of any of claims 17-25, wherein Y is -O-, -N(R<sub>a</sub>)- or -P-.
27. The method of any of claims 17-26, wherein Y is S.
28. The method of any of claims 17-27, wherein -N(R<sub>a</sub>)(R<sub>b</sub>) is amino.
29. The method of any of claims 17-28, wherein halo is Br or F.
30. The method of any of claims 17-29, wherein -N(R<sub>a</sub>)(R<sub>b</sub>) is pyrrolidino, piperidino or morpholino.
31. The method any of claims 1-4, wherein the antagonist of the IL-4 or IL-13 signal transduction pathway is a compound of formula (II):



(II)


wherein R<sup>1</sup>, R<sup>2</sup>, R<sup>3</sup> and R<sup>4</sup> are independently hydrogen, halo, hydroxy, cyano, , -N(R<sub>a</sub>)(R<sub>b</sub>), -S(R<sub>a</sub>), -NO<sub>2</sub>, (C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy, (C<sub>2</sub>-C<sub>6</sub>)alkynyl, (C<sub>2</sub>-C<sub>6</sub>)alkenyl, (C<sub>2</sub>-C<sub>7</sub>)alkanoyl, (C<sub>2</sub>-C<sub>7</sub>)alkanoyloxy, or (C<sub>3</sub>-C<sub>7</sub>)cycloalkyl or R<sup>1</sup> and R<sup>2</sup> taken together are benzo, optionally substituted by R<sup>1</sup>, (C<sub>3</sub>-C<sub>5</sub>)alkylene or methylene dioxy; wherein R<sub>a</sub> and R<sub>b</sub> are each independently hydrogen, (C<sub>1</sub>-C<sub>3</sub>)alkyl, (C<sub>2</sub>-C<sub>4</sub>)alkanoyl, phenyl, benzyl, or phenethyl; or R<sub>a</sub> and R<sub>b</sub> together with the nitrogen to which they are attached are a 5-6 membered heterocyclic ring, or R<sup>1</sup> and R<sup>4</sup> together with the atoms to which they are attached are benzo, (C<sub>3</sub>-C<sub>5</sub>)alkylidene or methylenedioxy;

Ar is aryl, heteroaryl, or a 5-6 membered heterocyclic ring, optionally comprising 1-3 N(R<sub>a</sub>), nonperoxide O or S atoms, optionally substituted with 1-5 groups where the groups are selected from halo,

CF<sub>3</sub>, hydroxy, CN, -N(R<sub>a</sub>)(R<sub>b</sub>), (C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy, (C<sub>2</sub>-C<sub>7</sub>)-alkanoyl, (C<sub>2</sub>-C<sub>7</sub>)alkanoyloxy, (C<sub>3</sub>-C<sub>7</sub>)cycloalkyl, (C<sub>2</sub>-C<sub>6</sub>)alkenyl, and phenyl;


or a pharmaceutically acceptable salt thereof.

32. The method of claim 31, wherein the heterocyclic ring, is pyrrolidino, piperidino or morpholino.
33. The method of any of claims 31-32, wherein the aryl, heteroaryl or heterocyclic group is substituted with 1-2 groups.
34. The method of any of claims 31-33, wherein N(R<sub>a</sub>)(R<sub>b</sub>) is pyrrolidino, piperidino or morpholino.
35. The method of any of claims 31-34, wherein the compound of formula (II) has formula (IIa) or (IIb):



or a pharmaceutically acceptable salt thereof.

36. The method any of claims 1-4, wherein the antagonist of the IL-4 or IL-13 signal transduction pathway is a compound of formula (III):




wherein  $R^1$ ,  $R^2$ , and  $R^4$  are independently hydrogen, halo, hydroxy, cyano,  $N(R_a)(R_b)$ ,  $S(R_a)$ ,  $NO_2$ ,  $(C_1-C_6)$ alkyl,  $(C_1-C_6)$ alkoxy,  $(C_2-C_6)$ alkynyl,  $(C_2-C_6)$ alkenyl,  $(C_2-C_7)$ alkanoyl,  $(C_2-C_7)$ alkanoyloxy, or  $(C_3-C_7)$ cycloalkyl or  $R^1$  and  $R^2$  taken together are benzo, optionally substituted by  $R^1$ ,  $(C_3-C_5)$ alkylene or methylene dioxy; wherein  $R_a$  and  $R_b$  are each independently hydrogen,  $(C_1-C_3)$ alkyl,  $(C_2-C_4)$ alkanoyl, phenyl, benzyl, or phenethyl; or  $R_a$  and  $R_b$  together with the nitrogen to which they are attached are a 5-6 membered heterocyclic ring, or  $R^1$  and  $R^4$  together with the atoms to which they are attached are benzo,  $(C_3-C_5)$ alkylidene or methylenedioxy;

Ar is aryl, heteroaryl, or a 5-6 membered heterocyclic ring, optionally comprising 1-3  $N(R_a)$ , nonperoxide O or S atoms, optionally substituted with 1-5 groups where the groups are selected from halo,  $CF_3$ , hydroxy, CN,  $-N(R_a)(R_b)$ ,  $(C_1-C_6)$ alkyl,  $(C_1-C_6)$ alkoxy,  $(C_2-C_7)$ alkanoyl,  $(C_2-C_7)$ alkanoyloxy,  $(C_3-C_7)$ cycloalkyl,  $(C_2-C_6)$ alkenyl, and phenyl;

or a pharmaceutically acceptable salt thereof.

37. The method of claim 36, wherein the heterocyclic ring, is pyrrolidino, piperidino or morpholino.
38. The method of claims 36 or 37, wherein the aryl, heteroaryl or heterocyclic group is substituted with 1-2 groups.
39. The method of any of claims 1-4, wherein the antagonist of the IL-4 or IL-13 signal transduction pathway is a compound of formula (IV):



(IV)




wherein  $R^1$ ,  $R^2$  and  $R^4$  are independently hydrogen, halo, hydroxy, cyano,  $N(R_a)(R_b)$ ,  $S(R_a)$ ,  $NO_2$ ,  $(C_1-C_6)$ alkyl,  $(C_1-C_6)$ alkoxy,  $(C_2-C_6)$ alkynyl,  $(C_2-C_6)$ alkenyl,  $(C_2-C_7)$ alkanoyl,  $(C_2-C_7)$ alkanoyloxy, or  $(C_3-C_7)$ cycloalkyl or  $R^1$  and  $R^2$  taken together are benzo, optionally substituted by  $R^1$ ,  $(C_3-C_7)$ cycloalkyl or methylene dioxy; wherein  $R_a$  and  $R_b$  are each independently hydrogen,  $(C_1-C_3)$ alkyl,  $(C_2-C_4)$ alkanoyl, phenyl, benzyl, or phenethyl; or  $R_a$  and  $R_b$  together with the nitrogen to which they are attached are a 5-6 membered heterocyclic ring;

Ar is aryl, heteroaryl, or a 5-6 membered heterocyclic ring, optionally comprising 1-3  $N(R_a)$ , nonperoxide O or S atoms, optionally substituted with 1-5 groups where the groups are selected from halo,  $CF_3$ , hydroxy, CN,  $-N(R_a)(R_b)$ ,  $(C_1-C_6)$ alkyl,  $(C_1-C_6)$ alkoxy,  $(C_2-C_7)$ alkanoyl,  $(C_2-C_7)$ alkanoyloxy,  $(C_3-C_7)$ cycloalkyl,  $(C_2-C_6)$ alkenyl, and phenyl;

or a pharmaceutically acceptable salt thereof.

40. The method of claim 39, wherein the heterocyclic ring, is pyrrolidino, piperidino or morpholino.
41. The method of claims 39 or 40, wherein the aryl, heteroaryl or heterocyclic group is substituted with 1-2 groups.
42. The method any of claims 1-4, wherein the antagonist of the IL-4 or IL-13 signal transduction pathway is a compound of formula (V):



(V)


wherein  $R^1$ ,  $R^2$ ,  $R^3$  and  $R^4$  are independently hydrogen, halo, hydroxy, cyano,  $N(R_a)(R_b)$ ,  $S(R_a)$ ,  $NO_2$ ,  $(C_1-C_6)$ alkyl,  $(C_1-C_6)$ alkoxy,  $(C_2-C_6)$ -

alkynyl, (C<sub>2</sub>-C<sub>6</sub>)alkenyl, (C<sub>2</sub>-C<sub>7</sub>)alkanoyl, (C<sub>2</sub>-C<sub>7</sub>)alkanoyloxy, or (C<sub>3</sub>-C<sub>7</sub>)cycloalkyl or R<sup>1</sup> and R<sup>2</sup> taken together are benzo, optionally substituted by R<sup>1</sup>, (C<sub>3</sub>-C<sub>5</sub>)alkylene or methylene dioxy; wherein R<sub>a</sub> and R<sub>b</sub> are each independently hydrogen, (C<sub>1</sub>-C<sub>3</sub>)alkyl, (C<sub>2</sub>-C<sub>4</sub>)alkanoyl, phenyl, benzyl, or phenethyl; or R<sub>a</sub> and R<sub>b</sub> together with the nitrogen to which they are attached are a 5-6 membered heterocyclic ring;

Ar is aryl, heteroaryl, or a 5-6 membered heterocyclic ring, optionally comprising 1-3 N(R<sub>a</sub>), nonperoxide O or S atoms, optionally substituted with 1-5 groups where the groups are selected from halo, CF<sub>3</sub>, hydroxy, CN, -N(R<sub>a</sub>)(R<sub>b</sub>), (C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy, (C<sub>2</sub>-C<sub>7</sub>)alkanoyl, (C<sub>2</sub>-C<sub>7</sub>)alkanoyloxy, (C<sub>3</sub>-C<sub>7</sub>)cycloalkyl, (C<sub>2</sub>-C<sub>6</sub>)alkenyl, and phenyl;

or a pharmaceutically acceptable salt thereof.

43. The method of claim 42, wherein the heterocyclic ring, is pyrrolidino, piperidino or morpholino.
44. The method of claims 42 and 43, wherein the aryl, heteroaryl or heterocyclic group is substituted with 1-2 groups.
45. The method any of claims 1-45, wherein the interferon is administered with of a compound of formula (I), (II), (III), (IV), (V) or a mixture thereof.
46. The use of a compound of formula (I):



(I)

wherein R<sup>1</sup>, R<sup>2</sup> and R<sup>3</sup> are independently hydrogen, halo, hydroxy, cyano, N(R<sub>a</sub>)(R<sub>b</sub>), S(R<sub>a</sub>), NO<sub>2</sub>, (C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy, (C<sub>2</sub>-C<sub>6</sub>)alkynyl, (C<sub>2</sub>-C<sub>6</sub>)alkenyl, (C<sub>2</sub>-C<sub>7</sub>)alkanoyl, (C<sub>2</sub>-C<sub>7</sub>)alkanoyloxy, or

(C<sub>3</sub>-C<sub>7</sub>)cycloalkyl or R<sup>1</sup> and R<sup>2</sup> taken together are benzo, optionally substituted by R<sup>1</sup>, (C<sub>3</sub>-C<sub>7</sub>)cycloalkyl or methylene dioxy; wherein R<sub>a</sub> and R<sub>b</sub> are each independently hydrogen, (C<sub>1</sub>-C<sub>3</sub>)alkyl, (C<sub>2</sub>-C<sub>4</sub>)alkanoyl, phenyl, benzyl, or phenethyl; or R<sub>a</sub> and R<sub>b</sub> together with the nitrogen to which they are attached are a 5-6 membered heterocyclic ring;

Ar is aryl or heteroaryl, optionally substituted with 1-5 CF<sub>3</sub>, hydroxy, CN, -N(R<sub>a</sub>)(R<sub>b</sub>), (C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>2</sub>-C<sub>7</sub>)alkanoyl, (C<sub>2</sub>-C<sub>7</sub>)alkanoyloxy, (C<sub>3</sub>-C<sub>7</sub>)cycloalkyl, (C<sub>2</sub>-C<sub>6</sub>)alkenyl, or phenyl groups; and

Y is oxy (-O-), S(O)<sub>0-2</sub>, Se, C(R<sup>1</sup>)(R<sup>3</sup>), N(R<sub>a</sub>), or -P-;


or a pharmaceutically acceptable salt thereof, to prepare a medicament effective to enhance the efficacy of interferon treatment.

47. The method of claim 46, wherein the heterocyclic ring, is pyrrolidino, piperidino or morpholino.
48. The method of claims 46 or 47, wherein the aryl, heteroaryl or heterocyclic group is substituted with 1-2 groups.
49. The use of any of claims 46-48, wherein R<sup>1</sup> or R<sup>2</sup> are independently hydroxy, cyano, -N(R<sub>a</sub>)(R<sub>b</sub>), S(R<sub>a</sub>), NO<sub>2</sub>, (C<sub>2</sub>-C<sub>7</sub>)alkanoyl, or (C<sub>2</sub>-C<sub>7</sub>)alkanoyloxy;
 

Ar is heteroaryl or phenyl substituted with cyano, (C<sub>2</sub>-C<sub>7</sub>)alkanoyl, (C<sub>2</sub>-C<sub>7</sub>)alkanoyloxy, (C<sub>2</sub>-C<sub>7</sub>)cycloalkyl or (C<sub>2</sub>-C<sub>6</sub>)alkenyl; and

Y is Se, SO, SO<sub>2</sub>, C(R<sub>1</sub>)(R<sub>3</sub>) or P.
50. The use of any of claims 46-49, wherein R<sup>1</sup> and R<sup>2</sup> together is butylene or benzo.
51. The use of any of claims 46-50, wherein R<sup>3</sup> is H.
52. The use of any of claims 46-51, wherein Y is oxy (-O-), S(O)<sub>0-2</sub>, C(R<sup>1</sup>)(R<sup>3</sup>), NR<sup>1</sup>, or -P-.
53. The use of any of claims 46-52, wherein Y is O, N(R<sub>a</sub>) or P.

54. The use of any of claims 46-53, wherein Ar is phenyl, 4-pyridyl or 2-thienyl.
55. The use of any of claims 46-54, wherein Ar is heteroaryl or phenyl substituted with CN, (C<sub>2</sub>-C<sub>7</sub>)alkanoyl, (C<sub>2</sub>-C<sub>7</sub>)alkanolyoxy, (C<sub>2</sub>-C<sub>7</sub>)cycloalkyl or (C<sub>2</sub>-C<sub>6</sub>)alkenyl.
56. The use of any of claims 46-55, wherein Ar is a 5-6 membered heterocyclic ring, comprising 1-3 N(R<sub>a</sub>), nonperoxide O or S atoms.
57. The use of any of claims 46-56, wherein Ar is pyrrolidino, piperidino or morpholino.
58. The use of any of claims 46-57, wherein Y is S.
59. The use of any of claims 46-58, wherein N(R<sub>a</sub>)(R<sub>b</sub>) is amino.
60. The method of any of claims 46-59, wherein halo is Br or F.
61. The use of any of claims 46-60, wherein N(R<sub>a</sub>)(R<sub>b</sub>) is pyrrolidino, piperidino or morpholino.
62. The use of a compound of formula (II):



(II)


wherein R<sup>1</sup>, R<sup>2</sup>, R<sup>3</sup> and R<sup>4</sup> are independently hydrogen, halo, hydroxy, cyano, N(R<sub>a</sub>)(R<sub>b</sub>), S(R<sub>a</sub>), NO<sub>2</sub>, (C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy, (C<sub>2</sub>-C<sub>6</sub>)-alkynyl, (C<sub>2</sub>-C<sub>6</sub>)alkenyl, (C<sub>2</sub>-C<sub>7</sub>)alkanoyl, (C<sub>2</sub>-C<sub>7</sub>)alkanoyloxy, or (C<sub>3</sub>-C<sub>7</sub>)cycloalkyl or R<sup>1</sup> and R<sup>2</sup> taken together are benzo, optionally substituted by R<sup>1</sup>, (C<sub>3</sub>-C<sub>5</sub>)alkylene or methylene dioxy; wherein R<sub>a</sub> and R<sub>b</sub> are each independently hydrogen, (C<sub>1</sub>-C<sub>3</sub>)alkyl, (C<sub>2</sub>-C<sub>4</sub>)alkanoyl,

phenyl, benzyl, or phenethyl; or  $R_a$  and  $R_b$  together with the nitrogen to which they are attached are a 5-6 membered heterocyclic ring, or  $R^1$  and  $R^4$  together with the atoms to which they are attached are benzo,  $(C_3-C_5)$ alkylidene or methylenedioxy;


Ar is aryl, heteroaryl, or a 5-6 membered heterocyclic ring, optionally comprising 1-3  $N(R_a)$ , nonperoxide O or S atoms, optionally substituted with 1-5 groups where the groups are selected from halo,  $CF_3$ , hydroxy, CN,  $-N(R_a)(R_b)$ ,  $(C_1-C_6)$ alkyl,  $(C_1-C_6)$ alkoxy,  $(C_2-C_7)$ -alkanoyl,  $(C_2-C_7)$ alkanoyloxy,  $(C_3-C_7)$ cycloalkyl,  $(C_2-C_6)$ alkenyl, and phenyl; or

or a pharmaceutically acceptable salt thereof, to prepare a medicament effective to enhance the efficacy of interferon treatment.

63. The method of claim 62, wherein the heterocyclic ring, is pyrrolidino, piperidino or morpholino.
64. The method of claim 62 or 63, wherein the aryl, heteroaryl or heterocyclic group is substituted with 1-2 groups.
65. The use of any of claims 62-64, wherein one of  $R^1$ ,  $R^2$ ,  $R^3$  and  $R^4$  is  $N(R_a)(R_b)$  and with the nitrogen to which they are attached is pyrrolidino, piperidino or morpholino.
66. The use of any of claims 62-65, wherein the compound of formula (II) has formula (IIa) or (IIb):




(IIa) ;



(IIb)

or a pharmaceutically acceptable salt thereof.

67. The use of a compound having formula (III):



(III)


wherein  $R^1$ ,  $R^2$ , and  $R^4$  are independently hydrogen, halo, hydroxy, cyano,  $N(R_a)(R_b)$ ,  $S(R_a)$ ,  $NO_2$ ,  $(C_1-C_6)$ alkyl,  $(C_1-C_6)$ alkoxy,  $(C_2-C_6)$ alkynyl,  $(C_2-C_6)$ alkenyl,  $(C_2-C_7)$ alkanoyl,  $(C_2-C_7)$ alkanoyloxy, or  $(C_3-C_7)$ cycloalkyl or  $R^1$  and  $R^2$  taken together are benzo, optionally substituted by  $R^1$ ,  $(C_3-C_5)$ alkylene or methylene dioxy; wherein  $R_a$  and  $R_b$  are each independently hydrogen,  $(C_1-C_3)$ alkyl,  $(C_2-C_4)$ alkanoyl, phenyl, benzyl, or phenethyl; or  $R_a$  and  $R_b$  together with the nitrogen to which they are attached are a 5-6 membered heterocyclic ring, or  $R^1$  and  $R^4$  together with the atoms to which they are attached are benzo,  $(C_3-C_5)$ alkylidene or methylenedioxy;

Ar is aryl, heteroaryl, or a 5-6 membered heterocyclic ring, optionally comprising 1-3  $N(R_a)$ , nonperoxide O or S atoms, optionally substituted with 1-5 groups where the groups are selected from halo,  $CF_3$ , hydroxy, CN,  $-N(R_a)(R_b)$ ,  $(C_1-C_6)$ alkyl,  $(C_1-C_6)$ alkoxy,  $(C_2-C_7)$ -alkanoyl,  $(C_2-C_7)$ alkanoyloxy,  $(C_3-C_7)$ cycloalkyl,  $(C_2-C_6)$ alkenyl, and phenyl; or

a pharmaceutically acceptable salt thereof, to prepare a medicament effective to enhance the efficacy of interferon treatment.

68. The method of claim 67, wherein the heterocyclic ring, is pyrrolidino, piperidino or morpholino.
69. The method of claim 67 or 68, wherein the aryl, heteroaryl or heterocyclic group is substituted with 1-2 groups.

70. The use of a compound having formula (IV):



(IV)


wherein R<sup>1</sup>, R<sup>2</sup> and R<sup>4</sup> are independently hydrogen, halo, hydroxy, cyano, N(R<sub>a</sub>)(R<sub>b</sub>), S(R<sub>a</sub>), NO<sub>2</sub>, (C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy, (C<sub>2</sub>-C<sub>6</sub>)alkynyl, (C<sub>2</sub>-C<sub>6</sub>)alkenyl, (C<sub>2</sub>-C<sub>7</sub>)alkanoyl, (C<sub>2</sub>-C<sub>7</sub>)alkanoyloxy, or (C<sub>3</sub>-C<sub>7</sub>)cycloalkyl or R<sup>1</sup> and R<sup>2</sup> taken together are benzo, optionally substituted by R<sup>1</sup>, (C<sub>3</sub>-C<sub>5</sub>)alkylene or methylene dioxy; wherein R<sub>a</sub> and R<sub>b</sub> are each independently hydrogen, (C<sub>1</sub>-C<sub>3</sub>)alkyl, (C<sub>2</sub>-C<sub>4</sub>)alkanoyl, phenyl, benzyl, or phenethyl; or R<sub>a</sub> and R<sub>b</sub> together with the nitrogen to which they are attached are a 5-6 membered heterocyclic ring;

Ar is aryl, heteroaryl, or a 5-6 membered heterocyclic ring, optionally comprising 1-3 N(R<sub>a</sub>), nonperoxide O or S atoms, optionally substituted with 1-5 groups where the groups are selected from halo, CF<sub>3</sub>, hydroxy, CN, -N(R<sub>a</sub>)(R<sub>b</sub>), (C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy, (C<sub>2</sub>-C<sub>7</sub>)alkanoyl, (C<sub>2</sub>-C<sub>7</sub>)alkanoyloxy, (C<sub>3</sub>-C<sub>7</sub>)cycloalkyl, (C<sub>2</sub>-C<sub>6</sub>)alkenyl, and phenyl; or

a pharmaceutically acceptable salt thereof, to prepare a medicament effective to enhance the efficacy of interferon treatment.

71. The method of claim 70, wherein the heterocyclic ring, is pyrrolidino, piperidino or morpholino.
72. The method of claim 70 or 71, wherein the aryl, heteroaryl or heterocyclic group is substituted with 1-2 groups.

73. The use of a compound having formula (V):



(V)

wherein  $R^1$ ,  $R^2$ ,  $R^3$  and  $R^4$  are independently hydrogen, halo, hydroxy, cyano,  $N(R_a)(R_b)$ ,  $S(R_a)$ ,  $NO_2$ ,  $(C_1-C_6)$ alkyl,  $(C_1-C_6)$ alkoxy,  $(C_2-C_6)$ alkynyl,  $(C_2-C_6)$ alkenyl,  $(C_2-C_7)$ alkanoyl,  $(C_2-C_7)$ alkanoyloxy, or  $(C_3-C_7)$ cycloalkyl or  $R^1$  and  $R^2$  taken together are benzo, optionally substituted by  $R^1$ ,  $(C_3-C_5)$ alkylene or methylene dioxy; wherein  $R_a$  and  $R_b$  are each independently hydrogen,  $(C_1-C_3)$ alkyl,  $(C_2-C_4)$ alkanoyl, phenyl, benzyl, or phenethyl; or  $R_a$  and  $R_b$  together with the nitrogen to which they are attached are a 5-6 membered heterocyclic ring, or  $R^1$  and  $R^4$  together with the atoms to which they are attached are benzo,  $(C_3-C_5)$ alkylidene or methylenedioxy;

Ar is aryl, heteroaryl, or a 5-6 membered heterocyclic ring, optionally comprising 1-3  $N(R_a)$ , nonperoxide O or S atoms, optionally substituted with 1-5 groups where the groups are selected from halo,  $CF_3$ , hydroxy, CN,  $-N(R_a)(R_b)$ ,  $(C_1-C_6)$ alkyl,  $(C_1-C_6)$ alkoxy,  $(C_2-C_7)$ -alkanoyl,  $(C_2-C_7)$ alkanoyloxy,  $(C_3-C_7)$ cycloalkyl,  $(C_2-C_6)$ alkenyl, and phenyl; or

a pharmaceutically acceptable salt thereof, to prepare a medicament effective to enhance the efficacy of interferon treatment.

74. The method of claim 73, wherein the heterocyclic ring, is pyrrolidino, piperidino or morpholino.
75. The method of claim 73 or 74, wherein the aryl, heteroaryl or heterocyclic group is substituted with 1-2 groups.






Fig.1A




Fig.1B

## CLL#1: PROTECTION AGAINST SPONTANEOUS APOPTOSIS




Fig.2A

## CLL#2: PROTECTION AGAINST FLUDARABINE-INDUCED APOPTOSIS




Fig.2B

## BEAS-2B SCREENING ASSAY




Fig.3




Fig.4A




Fig.4B




Fig.5A




Fig.5B




Fig.5C




Fig.5D




Fig.5E




Fig.5F




Fig.5G




Fig.5H




Fig.5I




Fig.5J




Fig.5K




Fig.5L




Fig.5M




Fig.5N




Fig.5O

Fig. 6
















Fig 7

## Inhibition of STAT-6




The BEAS-2B human airway epithelial cells were transiently transfected with the human 12/15-lipoxygenase promoter/luciferase reporter plasmid with and without a STAT6 expressing plasmid. Cells were then incubated with the indicated compound at 10  $\mu$ M for 1 hour, followed by IL-4 (10 ng/ml). After 16 hours, luciferase was measured using a chemiluminometer. The STAT-6 induction was normalized using the  $\beta$ -gal results as "background". The viability of the treated cells was visually verified at the end of the incubation, and found to be >95%.

BEST AVAILABLE COPY


## Fig. 8

# Measurement of Anti-STAT-6 Activity




The BEAS-2B human airway epithelial cells were transiently transfected with the human 12/15-lipoxygenase promoter/luciferase reporter gene. Cells were then incubated with the compound # 8 at the indicated concentrations for 1 hour, followed by IL-4 (10 ng/ml). After 16 hours, luciferase was measured using a chemiluminometer. The STAT-6 induction was normalized using the  $\beta$ -gal results as "background". Results are the mean of triplicate measurements, all SD are lower than 8%.

Fig. 9  
Measurement of Anti-STAT-6 Activity




The inhibitory activity of the compounds was tested at a fixed concentration of 10  $\mu$ M in IL-4 stimulated BEAS-2B human airway using the 12/15-lipoxygenase promoter/luciferase described in the previous figures.

Fig.10  
Functional Measurement of Anti-STAT-6 Activity




The effect of compounds 8, 12 and 28 on the IL-4-induced expression of the protein 12/15 lipooxygenase (12/15-LOX) was measured in primary human monocytes. Cells were pre-incubated for 1 hour in presence of 10  $\mu$ M of the compound, then exposed to 10 ng/ml of IL-4. The expression of 12/15-LOX was quantified by immunoblotting using a specific monoclonal antibody and compared to a loading control.

**Fig.11**  
Interferon Regulation Using IBT (Compound 8)



The B cells RAMOS were incubated as indicated with leukocyte-purified interferons (IFN,  $10^4$  U/ml), IL-4 (10 ng/ml) and compound #3 (10  $\mu$ M) for 48 hrs. The surface expression of the Fas receptor (FasR or CD95) was then measured by flow cytometry using a PE-conjugated monoclonal antibody. The percentage of CD95+ cells is indicated in each panel.

Fig 12

**BEST AVAILABLE COPY**

(19) World Intellectual Property  
Organization  
International Bureau



(43) International Publication Date  
17 April 2003 (17.04.2003)

PCT

(10) International Publication Number  
**WO 2003/031587 A3**

(51) International Patent Classification<sup>7</sup>: **A61K 31/519**,  
31/425, 31/42, 31/415

(21) International Application Number:  
PCT/US2002/032503

(22) International Filing Date: 9 October 2002 (09.10.2002)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
60/328,162 9 October 2001 (09.10.2001) US  
60/328,689 10 October 2001 (10.10.2001) US

(71) Applicants and

(72) Inventors: **CARSON, Dennis, A.** [US/US]; 14824 Vista del Oceano, Del Mar, CA 92014 (US). **COTTAM, Howard, B.** [US/US]; 3144 Willow Creek Place, Escondido, CA 92027 (US). **LEONI, Lorenzo, M.** [CH/US]; 3908 Via Tranquilo, San Diego, CA 92122 (US). **BARCHECHATH, Sylvie** [US/US]; 9321 Discovery Way, La Jolla, CA 92037 (US).


(74) Agents: **STEFFEY, Charles, E.** et al.; Schwegman, Lundberg, Woessner & Kluth, P.O. Box 2938, Minneapolis, MN 55402 (US).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),

[Continued on next page]

(54) Title: USE OF STAT-6 INHIBITORS AS THERAPEUTIC AGENTS



(57) Abstract: The present invention provides A therapeutic method to enhance the efficacy of interferon treatment comprising administering to a mammal subject to interferon treatment a compound which is an antagonist of the IL-4 or IL-13 signal transduction pathway in an amount effective to enhance said efficacy. The method includes treatment of diseases such as cancer, proliferative fibrotic diseases, viral diseases, or autoimmune diseases. The invention also includes the use of chemotherapeutic agents, radiation or other treatments in conjunction with the method of the invention.



European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

(88) Date of publication of the international search report:  
19 February 2004

**Published:**

- with international search report
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments

*For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*

**BEST AVAILABLE COPY**



## INTERNATIONAL SEARCH REPORT

International application No.

PCT/US02/32503

## A. CLASSIFICATION OF SUBJECT MATTER

IPC(7) : A61K 31/519,31/425,31/42,31/415

US CL : 514/256,257,258,259,260,365,367,368,370,374,375,377,385,396,397

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 514/256,257,258,259,260,365,367,368,370,374,375,377,385,396,397

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)  
CAS ONLINE, MEDLINE search terms: interferon, IL-3, IL-4, cancer

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	US 4,762,705 A (RUBIN) 09 August 1988 (09.08.1988), the entire document, particularly, the abstract, and the claims	1-3, 46-48, 62-64, 67-73
Y	JP 11-106340 (INUOE et al) 20 April 1999 (20.04.1999), the abstract, and the claims.	1-3, 46-48, 62-64, 67-73
Y	WO 97/42192 A1 (BASF AKTIENGESELLSCHAFT) 13 November 1997 (13.11.1997), the abstract.	62-64
Y	US 4,567,263 A (EICKEN et al) 28 January 1986 (28.01.1986), the abstract and the claims.	67-69
Y	US 4,537,889 A (SPITZER) 27 August 1985 (27.08.1985), abstract, and the claims.	73
Y	FRANCIS et al, 'Anxiolytic properties of certain annelated [1,2,4] triazol[1,5-c]pyrimidin-5(6H)-ones,' J. Med. Chem. 1991, Vol. 34, pages 2899-2906, see the entire document.	70-72

☐ Further documents are listed in the continuation of Box C.☐ See patent family annex.

## \* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier application or patent published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T"

later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X"

document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y"

document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&amp;"

document member of the same patent family

Date of the actual completion of the international search

25 November 2003 (25.11.2003)

Date of mailing of the international search report

15 DEC 2003

Name and mailing address of the ISA/US

Mail Stop PCT, Attn: ISA/US

Commissioner for Patents

P.O. Box 1450

Alexandria, Virginia 22313-1450

Facsimile No. (703)305-3230

Authorized officer

Shengjun Wang

Telephone No. (703) 308-1235

Form PCT/ISA/210 (second sheet) (July 1998)

BEST AVAILABLE COPY

# INTERNATIONAL SEARCH REPORT

International application No.

PCT/US02/32503

## Box I Observations where certain claims were found unsearchable (Continuation of Item 1 of first sheet)

This international report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claim Nos.:  
because they relate to subject matter not required to be searched by this Authority, namely:
2. ☐ Claim Nos.:  
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
3. ☒ Claim Nos.: 4-45, 49-61, 65, 66  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

## Box II Observations where unity of invention is lacking (Continuation of Item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

☐  
☐

The additional search fees were accompanied by the applicant's protest.  
No protest accompanied the payment of additional search fees.